



Critical Care Skills Checklist

This profile helps us place you in an assignment suited to your skill level and expertise. This checklist is intended for Critical Care nurses with (1) or more years of experience in their discipline and specialty.

First Name: _____ Last Name: _____

Date: _____ S.S. Number: _____

Please indicate your current level of experience. There are no right/wrong answers.

- 1.) No Experience, Theory Only
- 2.) Limited Experience
- 3.) Moderate Experience
- 4.) Experienced and Competent
- 5.) Experienced; Able to teach

I. CARDIOVASCULAR

- | | | | | | |
|---|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| a. Abnormal heart sounds / murmurs | 1 <input type="radio"/> | 2 <input type="radio"/> | 3 <input type="radio"/> | 4 <input type="radio"/> | 5 <input type="radio"/> |
| b. Auscultation (rate, rhythm) | 1 <input type="radio"/> | 2 <input type="radio"/> | 3 <input type="radio"/> | 4 <input type="radio"/> | 5 <input type="radio"/> |
| c. Pulses / circulation checks | 1 <input type="radio"/> | 2 <input type="radio"/> | 3 <input type="radio"/> | 4 <input type="radio"/> | 5 <input type="radio"/> |
| d. Use of Doppler | 1 <input type="radio"/> | 2 <input type="radio"/> | 3 <input type="radio"/> | 4 <input type="radio"/> | 5 <input type="radio"/> |
| e. Interpretation of cardiac enzymes and isoenzymes | 1 <input type="radio"/> | 2 <input type="radio"/> | 3 <input type="radio"/> | 4 <input type="radio"/> | 5 <input type="radio"/> |
| f. Interpretation of coagulation studies | 1 <input type="radio"/> | 2 <input type="radio"/> | 3 <input type="radio"/> | 4 <input type="radio"/> | 5 <input type="radio"/> |
| g. Arterial line insertion | 1 <input type="radio"/> | 2 <input type="radio"/> | 3 <input type="radio"/> | 4 <input type="radio"/> | 5 <input type="radio"/> |
| h. Central line insertion | 1 <input type="radio"/> | 2 <input type="radio"/> | 3 <input type="radio"/> | 4 <input type="radio"/> | 5 <input type="radio"/> |
| i. Open Chest Emergency | 1 <input type="radio"/> | 2 <input type="radio"/> | 3 <input type="radio"/> | 4 <input type="radio"/> | 5 <input type="radio"/> |
| j. PA catheter / Swan-Ganz insertion | 1 <input type="radio"/> | 2 <input type="radio"/> | 3 <input type="radio"/> | 4 <input type="radio"/> | 5 <input type="radio"/> |
| k. Pericardiocentesis | 1 <input type="radio"/> | 2 <input type="radio"/> | 3 <input type="radio"/> | 4 <input type="radio"/> | 5 <input type="radio"/> |

l. Automatic internal cardioverter defibrillator	1 ○	2 ○	3 ○	4 ○	5 ○
m. Cardioversion / Defibrillation	1 ○	2 ○	3 ○	4 ○	5 ○
n. Cardiac index	1 ○	2 ○	3 ○	4 ○	5 ○
o. Cardiac output	1 ○	2 ○	3 ○	4 ○	5 ○
p. CVP monitoring	1 ○	2 ○	3 ○	4 ○	5 ○
q. MAP monitoring	1 ○	2 ○	3 ○	4 ○	5 ○
r. P/A Swan-Ganz	1 ○	2 ○	3 ○	4 ○	5 ○
s. PVR	1 ○	2 ○	3 ○	4 ○	5 ○
t. Radial a-line	1 ○	2 ○	3 ○	4 ○	5 ○
u. SVO2 (mixed venous oxyhemoglobin)	1 ○	2 ○	3 ○	4 ○	5 ○
v. SVR (systemic vascular resistance)	1 ○	2 ○	3 ○	4 ○	5 ○
w. Arrhythmia interpretation	1 ○	2 ○	3 ○	4 ○	5 ○
x. Lead placement	1 ○	2 ○	3 ○	4 ○	5 ○
y. Rhythm strip assessment	1 ○	2 ○	3 ○	4 ○	5 ○
z. Set up, run, and interpretation of 12 lead EKG	1 ○	2 ○	3 ○	4 ○	5 ○
a. Temporary external pacing	1 ○	2 ○	3 ○	4 ○	5 ○
b. Temporary transvenous	1 ○	2 ○	3 ○	4 ○	5 ○
c. Temporary epicardial wires	1 ○	2 ○	3 ○	4 ○	5 ○
d. Permanent	1 ○	2 ○	3 ○	4 ○	5 ○
e. Abdominal aortic aneurysm repair	1 ○	2 ○	3 ○	4 ○	5 ○
f. Acute MI	1 ○	2 ○	3 ○	4 ○	5 ○
g. Cardiac arrest	1 ○	2 ○	3 ○	4 ○	5 ○
h. Cardiac tamponade	1 ○	2 ○	3 ○	4 ○	5 ○
i. CHF	1 ○	2 ○	3 ○	4 ○	5 ○
j. Heart transplant	1 ○	2 ○	3 ○	4 ○	5 ○
k. Immediate post open heart surgery	1 ○	2 ○	3 ○	4 ○	5 ○
l. Infective endocarditis	1 ○	2 ○	3 ○	4 ○	5 ○
m. Myocardial contusion	1 ○	2 ○	3 ○	4 ○	5 ○
n. Pericarditis	1 ○	2 ○	3 ○	4 ○	5 ○
o. PTCA (percutaneous transluminal coronary angioplasty)	1 ○	2 ○	3 ○	4 ○	5 ○
p. Post percutaneous balloon valvuloplasty	1 ○	2 ○	3 ○	4 ○	5 ○
q. Pre / post cardiac cath	1 ○	2 ○	3 ○	4 ○	5 ○

- | | | | | | |
|---|-----|-----|-----|-----|-----|
| r. Ventricular assist devices
(IABP, RVAD, LVAD) | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| s. Balloon Pumps | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| t. Atropine | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| u. Digoxin (Lanoxin) | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| v. Dobutamine (Dobutrex) | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| w. Dopamine (Intropin) | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| x. Epinephrine (Adrenalin) | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| y. Inocor (Amrinone) | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| z. Isuprel | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| a. Lidocaine (Xylocaine) | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| b. Nipride (Nitroprusside) | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| c. Nitroglycerine (Tridil) | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| d. Sodium Bicarbonate | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| e. TPA (Alteplase) | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |

II. PULMONARY

- | | | | | | |
|--|-----|-----|-----|-----|-----|
| a. Adventitious breath sounds | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| b. Rate and work of breathing | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| c. Interpretation of ABG's | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| d. Assist with intubation | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| e. Assist with extubation | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| f. Insertion of oral airway | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| g. chest physiotherapy | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| h. Incentive spirometer | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| i. Use of Pleurevac or other chest
drainage systems | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| j. Monitor pulse oximetry | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| k. Endotracheal tube suctioning | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| l. Oropharyngeal suctioning | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| m. Nasotracheal suctioning | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| n. Assist with bronchoscopy | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| o. Assist with chest tube insertion | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| p. Assist with emergency trach-
eostomy | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |

- | | | | | | |
|-----------------------------------|---|---|---|---|---|
| q. Assist with thoracentesis | 1 | 2 | 3 | 4 | 5 |
| r. Ambu bag techniques | 1 | 2 | 3 | 4 | 5 |
| s. External CPAP | 1 | 2 | 3 | 4 | 5 |
| t. High frequency jet ventilation | 1 | 2 | 3 | 4 | 5 |
| u. IMV | 1 | 2 | 3 | 4 | 5 |
| v. PEEP | 1 | 2 | 3 | 4 | 5 |
| w. Pressure support | 1 | 2 | 3 | 4 | 5 |
| x. Troubleshooting with vents | 1 | 2 | 3 | 4 | 5 |
| y. Weaning from ventilator | 1 | 2 | 3 | 4 | 5 |
| z. Acute pneumonia | 1 | 2 | 3 | 4 | 5 |
| a. ARDS | 1 | 2 | 3 | 4 | 5 |
| b. Chest trauma | 1 | 2 | 3 | 4 | 5 |
| c. COPD | 1 | 2 | 3 | 4 | 5 |
| d. Cor pulmonale | 1 | 2 | 3 | 4 | 5 |
| e. ECMO | 1 | 2 | 3 | 4 | 5 |
| f. Inhalation injuries | 1 | 2 | 3 | 4 | 5 |
| g. Lobectomy | 1 | 2 | 3 | 4 | 5 |
| h. Lung transplant | 1 | 2 | 3 | 4 | 5 |
| i. Near drowning | 1 | 2 | 3 | 4 | 5 |
| j. Pneumonectomy | 1 | 2 | 3 | 4 | 5 |
| k. Pulmonary edema | 1 | 2 | 3 | 4 | 5 |
| l. Pulmonary embolism | 1 | 2 | 3 | 4 | 5 |
| m. Thoracotomy | 1 | 2 | 3 | 4 | 5 |
| n. Tracheostomy | 1 | 2 | 3 | 4 | 5 |
| o. TB | 1 | 2 | 3 | 4 | 5 |
| p. Alupent (Metaprotenerol) | 1 | 2 | 3 | 4 | 5 |
| q. Corticosteroids | 1 | 2 | 3 | 4 | 5 |
| r. Ventolin (Albuterol) | 1 | 2 | 3 | 4 | 5 |

III. NEUROLOGY

- | | | | | | |
|------------------------------|---|---|---|---|---|
| a. Assess LOC | 1 | 2 | 3 | 4 | 5 |
| b. Use of Glasgow coma scale | 1 | 2 | 3 | 4 | 5 |
| c. Cranial nerve assessment | 1 | 2 | 3 | 4 | 5 |
| d. Seizure precautions | 1 | 2 | 3 | 4 | 5 |
| e. Spinal precautions | 1 | 2 | 3 | 4 | 5 |

- | | | | | | |
|---|---|---|---|---|---|
| f. Cervical traction | 1 | 2 | 3 | 4 | 5 |
| g. Halo traction | 1 | 2 | 3 | 4 | 5 |
| h. Assist with lumbar puncture | 1 | 2 | 3 | 4 | 5 |
| i. Hypo/hyperthermia regulation | 1 | 2 | 3 | 4 | 5 |
| j. ICP (intracranial pressure monitoring) | 1 | 2 | 3 | 4 | 5 |
| k. Nerve stimulators | 1 | 2 | 3 | 4 | 5 |
| l. Aneurysm precautions | 1 | 2 | 3 | 4 | 5 |
| m. Basal skull fracture | 1 | 2 | 3 | 4 | 5 |
| n. Closed head injury | 1 | 2 | 3 | 4 | 5 |
| o. Coma | 1 | 2 | 3 | 4 | 5 |
| p. CVA | 1 | 2 | 3 | 4 | 5 |
| q. DT's | 1 | 2 | 3 | 4 | 5 |
| r. Encephalitis | 1 | 2 | 3 | 4 | 5 |
| s. Externalized VP shunts | 1 | 2 | 3 | 4 | 5 |
| t. Increased ICP | 1 | 2 | 3 | 4 | 5 |
| u. Intracranial Hemorrhage | 1 | 2 | 3 | 4 | 5 |
| v. Laminectomy | 1 | 2 | 3 | 4 | 5 |
| w. Meningitis | 1 | 2 | 3 | 4 | 5 |
| x. Metastatic | 1 | 2 | 3 | 4 | 5 |
| y. Tumor/intracranial tumor
resection | 1 | 2 | 3 | 4 | 5 |
| z. MS (multiple sclerosis) | 1 | 2 | 3 | 4 | 5 |
| a. Post craniotomy | 1 | 2 | 3 | 4 | 5 |
| b. Seizure disorder | 1 | 2 | 3 | 4 | 5 |
| c. Spinal cord injury | 1 | 2 | 3 | 4 | 5 |
| d. Ventriculostomy | 1 | 2 | 3 | 4 | 5 |
| e. Decadron (Dexamethasone) | 1 | 2 | 3 | 4 | 5 |
| f. Dilantin (Phenytoin) | 1 | 2 | 3 | 4 | 5 |
| g. Epidural administration | 1 | 2 | 3 | 4 | 5 |
| h. Magnesium Sulfate | 1 | 2 | 3 | 4 | 5 |
| i. Phenobarbital | 1 | 2 | 3 | 4 | 5 |
| j. Valium (Diazepam) | 1 | 2 | 3 | 4 | 5 |

IV. GASTROINTESTINAL

- | | | | | | |
|--|---|---|---|---|---|
| a. Assessment of abdomen
(bowel sounds) | 1 | 2 | 3 | 4 | 5 |
|--|---|---|---|---|---|

- | | | | | | |
|--|---|---|---|---|---|
| b. Administration of tube feeding | 1 | 2 | 3 | 4 | 5 |
| c. Balloon tamponade (i.e.,
Sengstaken-Blakemore) | 1 | 2 | 3 | 4 | 5 |
| d. Feeding pump | 1 | 2 | 3 | 4 | 5 |
| e. Flexible feeding tube (i.e.,
Corpak, Dubhoff) | 1 | 2 | 3 | 4 | 5 |
| f. Gravit feeding | 1 | 2 | 3 | 4 | 5 |
| g. Iced saline lavage | 1 | 2 | 3 | 4 | 5 |
| h. Salem sump to suction | 1 | 2 | 3 | 4 | 5 |
| i. Gastrostomy tube | 1 | 2 | 3 | 4 | 5 |
| j. Jejunostomy tube (J-tube) | 1 | 2 | 3 | 4 | 5 |
| k. T-tube | 1 | 2 | 3 | 4 | 5 |
| l. Blunt trauma | 1 | 2 | 3 | 4 | 5 |
| m. Bowel obstruction | 1 | 2 | 3 | 4 | 5 |
| n. Colostomy | 1 | 2 | 3 | 4 | 5 |
| o. Esophageal bleeding | 1 | 2 | 3 | 4 | 5 |
| p. GI bleeding | 1 | 2 | 3 | 4 | 5 |
| q. GI surgery | 1 | 2 | 3 | 4 | 5 |
| r. Hepatitis | 1 | 2 | 3 | 4 | 5 |
| s. Ileostomy | 1 | 2 | 3 | 4 | 5 |
| t. IBD | 1 | 2 | 3 | 4 | 5 |
| u. Liver failure | 1 | 2 | 3 | 4 | 5 |
| v. Liver transplant | 1 | 2 | 3 | 4 | 5 |
| w. Pancreatitis | 1 | 2 | 3 | 4 | 5 |
| x. Paralytic ileus | 1 | 2 | 3 | 4 | 5 |
| y. Penetrating trauma | 1 | 2 | 3 | 4 | 5 |
| z. Aqua Mephyton (Vit. K) | 1 | 2 | 3 | 4 | 5 |
| a. Inderal (Propranolol) | 1 | 2 | 3 | 4 | 5 |
| b. Kayexelate | 1 | 2 | 3 | 4 | 5 |
| c. Lactulose (Cephulac) | 1 | 2 | 3 | 4 | 5 |
| d. Pitressin (Vasopressin) | 1 | 2 | 3 | 4 | 5 |
| e. TPN and lipid administration | 1 | 2 | 3 | 4 | 5 |
| f. PPN (peripheral parenteral nutrition) | 1 | 2 | 3 | 4 | 5 |

V. RENAL

- | | | | | | |
|-----------------------|---|---|---|---|---|
| a. Bladder irrigation | 1 | 2 | 3 | 4 | 5 |
|-----------------------|---|---|---|---|---|

- | | | | | | |
|--|-----|-----|-----|-----|-----|
| b. Insertion and care of Foley catheter | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| c. Acute renal failure | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| d. End stage renal failure | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| e. Hemodialysis | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| f. Nephrectomy | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| g. Nephrectomy tube placement | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| h. Peritoneal dialysis | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| i. Renal rejection syndrome | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| j. Renal transplant | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| k. Supra-pubic catheter | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| l. TURP | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| m. Urinary diversion (ileal conduit nephrostomy) | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| n. UTI | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |

VI. WOUND CARE

- | | | | | | |
|-------------------------------------|-----|-----|-----|-----|-----|
| a. Wound care irrigations | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| b. Sterile dressing changes | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| c. Burns | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| d. Pressure sores | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| e. Staged decubitus ulcers | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| f. Surgical wounds with drains (JP) | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| g. Traumatic wounds | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |

VII. IV THERAPY

- | | | | | | |
|------------------------------------|-----|-----|-----|-----|-----|
| a. IV peripheral starts | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| b. Drawing blood from central line | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| c. Drawing venous blood | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| d. Cryoprecipitate | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| e. Packed RBC | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| f. Plasma / albumin | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| g. Whole blood | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |

- | | | | | | |
|---|-----|-----|-----|-----|-----|
| h. IV calculations / titrate drip rates | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| i. Use of infusion pumps | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| j. Broviac | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| k. Groshong | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| l. Hickman | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| m. Portacath | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| n. Quinton | | | | | |

VII. OTHER NURSING RESPONSIBILITIES

- | | | | | | |
|--|-----|-----|-----|-----|-----|
| a. Charge nurse responsibilities | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| b. Isolation procedures/precautions | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| c. Use of restraints | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| d. Blood glucose monitoring | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| e. Insulin administration | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| f. Assess for signs and symptoms of insulin shock/diabetic coma | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| g. Oncology / chemotherapy administration | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| h. Collection and handling of lab specimens (blood, stool, urine, nasopharyngeal, wound) | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| i. Anaphylactic shock | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| j. Disseminated intravascular coagulation | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| k. Hypovolemic shock | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| l. MAST suit/rotating tourniquets | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| m. Organ/tissue donation | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |
| n. Septic shock | 1 ○ | 2 ○ | 3 ○ | 4 ○ | 5 ○ |

AGE SPECIFIC CRITERIA

Please indicate by circling each age group for which you have expertise in providing age-appropriate nursing care.

- A. Newborn/Neonate (birth – 30 days)
- B. Infant (30 days – 1 year)
- C. Toddler (1 – 3 years)
- D. Preschooler (3 – 5 years)

- E. School age children (5 – 12 years)
- F. Adolescents (12 – 18 years)
- G. Young Adults (18 – 39 years)
- H. Middle adults (39 – 64 years)
- I. Older adults (64 + years)

Able to adapt care to incorporate normal growth and development:
Please circle if you have experience in the following:

- A. Newborn/Neonate (birth – 30 days)
- B. Infant (30 days – 1 year)
- C. Toddler (1 – 3 years)
- D. Preschooler (3 – 5 years)
- E. School age children (5 – 12 years)
- F. Adolescents (12 – 18 years)
- G. Young Adults (18 – 39 years)
- H. Middle adults (39 – 64 years)
- I. Older adults (64 + years)

Able to adapt method/terminology of patient instruction to comprehension:
Please circle if you have experience in the following:

- A. Newborn/Neonate (birth – 30 days)
- B. Infant (30 days – 1 year)
- C. Toddler (1 – 3 years)
- D. Preschooler (3 – 5 years)
- E. School age children (5 – 12 years)
- F. Adolescents (12 – 18 years)
- G. Young Adults (18 – 39 years)
- H. Middle adults (39 – 64 years)
- I. Older adults (64 + years)

Can ensure a safe environment reflecting the special needs of various age groups:

- A. Newborn/Neonate (birth – 30 days)
- B. Infant (30 days – 1 year)
- C. Toddler (1 – 3 years)
- D. Preschooler (3 – 5 years)
- E. School age children (5 – 12 years)
- F. Adolescents (12 – 18 years)
- G. Young Adults (18 – 39 years)
- H. Middle adults (39 – 64 years)
- I. Older adults (64 + years)

Please read and agree to the statements below by signing and dating the bottom

I attest that the information and statements made in this checklist are true and accurate to the best of my knowledge. I am the individual completing this form. I understand that any falsification will be the basis for disqualification of employment. I authorize Signature Healthcare to verify the information I have provided and to contact past employers and references concerning my ability, character and employment record. I release all such persons from liability for furnishing said information. I hereby authorize Signature Healthcare to release this Critical Care Skills Checklist to client facilities in relation to consideration of employment as a Traveler, Contract or Per-diem worker with those facilities.

Signature

Date



3815 NW 109th Street Suite B
Urbandale, IA 50322
Human Resources Department
Christi J. Hokel, RN, BSN

Phone: 515.252.0000 or 1.800.518.1460

Fax us at:
515.276.5506